

## STUDY TIPS

# FA1.2: FACTORISATION: QUADRATICS

## Definition

The general form of a quadratic expression is:

$$ax^2 + bx + c \quad a \neq 0$$

where  $a$ ,  $b$ ,  $c$  are constants and  $x$  is the unknown.

## Expansion

Revision of expansion.

To expand an expression of the form  $(a + b)(c + d)$ , multiply each term in the first bracket by each term in the second bracket.

$$(x + 2)(x + 3) = x(x + 3) + 2(x + 3) = x^2 + 3x + 2x + 6$$

Simplify by combining like terms :

$$(x + 2)(x + 3) = x^2 + 5x + 6$$

## Factorisation

Factorisation is the reverse of expansion.

$x^2 + 5x + 6$  is expressed as the product of two factors,  $(x + 2)$  and  $(x + 3)$ .

$$\begin{array}{ccc}
 & \xrightarrow{\text{factorisation}} & \\
 x^2 + 5x + 6 = & (x + 2)(x + 3) & \\
 \xleftarrow{\text{expansion}} & & 
 \end{array}$$

Note that:

- multiplying the first term in each bracket gives the term ' $x^2$ ' in the expression
- multiplying the last term in each bracket gives the constant term, (+6), in the expression
- the coefficient of the ' $x$ ' term is the sum of last term in each bracket (+2 + 3 = +5).

In general

*constant 'c' positive*

$$x^2 + bx + c = (x + m)(x + n) \quad \text{and} \quad x^2 - bx + c = (x - m)(x - n)$$

*constant 'c' negative*

$$x^2 + bx - c = (x + m)(x - n) \quad m > n$$

$$x^2 - bx - c = (x + m)(x - n) \quad m < n \quad \text{where } m \text{ and } n \text{ are } \geq 0.$$

## Expressions with $a = 1$

If  $a = 1$  the expression becomes  $x^2 + bx + c$

To factorise:

Find two numbers that:

- multiply to 'c' and
- add to 'b'

Insert one number into each of the brackets  $(x \dots)(x \dots)$ .

The order is not important.

Note:

- If 'c' is positive, both numbers are positive or both numbers are negative.  
If b is positive, both numbers are positive.  
If b is negative, both numbers are negative.
- If c is negative, then one number is positive and the other negative;  
If b is positive, then the larger number is positive.  
If b is negative, then the larger number is negative.

## Examples

Factorise the following.

1.  $x^2 + 9x + 14$

$$\begin{aligned}x^2 + 9x + 14 &= (x \dots)(x \dots) \\ &= (x + 2)(x + 7)\end{aligned}$$

the first term in each bracket must be 'x'

Look for two numbers that

- multiply to +14
- add to +9

the numbers are +2 and +7

2.  $y^2 - 7y + 12$

$$\begin{aligned}y^2 - 7y + 12 &= (y \dots)(y \dots) \\ &= (y - 3)(y - 4)\end{aligned}$$

the first term in each bracket must be 'y'

Look for two numbers that

- multiply to +12.
- add to -7,

the numbers are -3 and -4

3.  $p^2 - 5p - 14$

$$\begin{aligned}p^2 - 5p - 14 &= (p \dots)(p \dots) \\ &= (p - 7)(p + 2)\end{aligned}$$

the first term in each bracket must be 'p'

Look for two numbers that

- multiply to -14
- add to -5,

(the larger number must be -ve)

numbers are -7 and +2

4.  $a^2 + 6a - 7$

$$\begin{aligned}a^2 + 6a - 7 &= (a \dots)(a \dots) \\ &= (a + 7)(a - 1)\end{aligned}$$

the first term in each bracket must be 'a'

Look for two numbers that

- multiply to -7
- add to +6

(the larger number must be +ve)

numbers are +7 and -1

Not all expressions of the form  $x^2 + bx + c$  can be factorised by this method. The factors may be of a more complicated form or there may be no real factors.

## Discriminant

For a quadratic equation of the form  $ax^2 + bx + c$  a method of checking for real factors is to calculate the value of  $(b^2 - 4ac)$  (called the *discriminant*).

If  $b^2 - 4ac < 0$  there are no real factors

## Example

Factorise (if possible)  $a^2 + 6a + 12$

In this case:

$$'b' = +6 \quad 'c' = +12$$

$$\therefore (b^2 - 4ac) = -12$$

The discriminant is negative therefore  $a^2 + 6a + 12$  has no real factors.

## Exercise

Factorise the following (if possible).

1.  $x^2 + 10x + 21$

4.  $m^2 - m - 72$

7.  $x^2 - 2x - 24$

10.  $n^2 + 6n - 16$

13.  $y^2 + 7y + 19$

2.  $z^2 + 11z + 18$

5.  $x^2 + 6x + 9$

8.  $y^2 - 10y + 16$

11.  $a^2 + 5a + 10$

14.  $x^2 + 16x + 39$

3.  $x^2 + 5x - 14$

6.  $a^2 - 15a + 44$

9.  $z^2 + 4z - 60$

12.  $s^2 + 2s - 48$

15.  $x^2 - 14x + 45$

## Answers

1.  $(x+7)(x+3)$

4.  $(m-9)(m+8)$

7.  $(x-6)(x+4)$

10.  $(n-2)(n+8)$

13. no real factors

2.  $(z+9)(z+2)$

5.  $(x+3)(x+3)$

8.  $(y-2)(y-8)$

11. no real factors

14.  $(x+13)(x+3)$

3.  $(x+7)(x-2)$

6.  $(a-11)(a-4)$

9.  $(z-6)(z+10)$

12.  $(s+8)(s-6)$

15.  $(x-9)(x-5)$

## Expressions with $a \neq 1$

Expressions of the type  $ax^2 + bx + c$  can be factorised using a technique similar to that used for expressions of the type  $x^2 + bx + c$ .

In this case the coefficient of  $x$ , in at least one bracket, will not equal 1.

Consider the following product.

$$\begin{aligned}(3x+2)(2x+1) &= 6x^2 + 3x + 4x + 2 \\ &= 6x^2 + 7x + 2\end{aligned}$$

- multiplying the first term in each bracket gives the ' $x^2$ ' term ( $6x^2$ ) in the expansion
- multiplying the last term in each bracket gives the constant term ( $+2$ ), in the expansion
- the coefficient of the ' $x$ ' term is the sum of the ' $x$ ' terms in the expansion. ( $+3x + 4x = +7x$ ).

## Examples

1. Factorise  $2x^2 + 7x + 6$
- $$\begin{aligned}2x^2 + 7x + 6 &= (2x\dots)(x\dots) \\ 2x^2 + 7x + 6 &= (2x+3)(x+2)\end{aligned}$$
- $2x \times x = 2x^2$
  - find factors of  $+6$   
middle term positive so  
 $+6$  and  $+1$  or  $+3$  and  $+2$
  - try all combinations.
  - the one that gives  $+7x$  when the brackets are expanded is  $(2x+3)$  and  $(x+2)$
2. Factorise  $2x^2 - x - 6$
- $$\begin{aligned}2x^2 - x - 6 &= (2x\dots)(x\dots) \\ 2x^2 - x - 6 &= (2x+3)(x-2)\end{aligned}$$
- $2x \times x = 2x^2$
  - find factors of  $-6$ .
  - $+6$  &  $-1$  or  $-6$  &  $+1$  or  $+3$  &  $-2$  or  $-3$  &  $+2$
  - try all combinations
  - the one that gives  $-x$  when the brackets are expanded is  $(2x+3)$  and  $(x-2)$
3. Factorise  $3a^2 - 16a + 5$
- $$\begin{aligned}3a^2 - 16a + 5 &= (3a\dots)(a\dots) \\ 3a^2 - 16a + 5 &= (a-5)(3a-1)\end{aligned}$$
- $3a \times a = 3a^2$
  - find factors of  $+5$
  - middle term negative so  $-5$  and  $-1$
  - try each combination
  - the one that gives  $-16a$  when the brackets are expanded is  $(a-5)$  and  $(3a-1)$
4. Factorise  $6y^2 + 16y - 6$
- $$\begin{aligned}6y^2 + 16y - 6 &= 2(3y^2 + 8y - 3) \\ &= 2(3y\dots)(y\dots) \\ 6y^2 + 16y - 6 &= 2(3y-1)(y+3)\end{aligned}$$
- Remove a common factor of 2.  
This reduces the number of possible combinations.
- $3y \times y = 3y^2$
  - find factors of  $-3$
  - possibilities are  $+3$  and  $-1$  or  $-3$  and  $+1$
  - try all combinations
  - the one that gives  $+8y$  when the brackets are expanded is  $(3y-1)$  and  $(y+3)$

To factorise a quadratic of the form  $ax^2 + bx + c$

- search each term in the expression for a common factor (**every term** must have this factor).
- remove this from the expression then place before the brackets.
- factorise, if possible, the resulting quadratic expression

This approach is good provided the number of combinations of factors of 'a' and 'c' is limited. For more complicated expressions of the type  $ax^2 + bx + c$  there are other methods of factorisation which may be more suitable.

### Exercise

Factorise the following

(1)  $5x^2 + 13x + 6$

(2)  $2x^2 + x - 15$

(3)  $3m^2 - m - 2$

(4)  $3y^2 - 10y + 8$

(5)  $2a^2 + 11a + 12$

(6)  $6x^2 - 11x + 5$

### Answers

(1)  $(5x + 3)(x + 2)$

(2)  $(2x - 5)(x + 3)$

(3)  $(3m + 2)(m - 1)$

(4)  $(3y - 4)(y - 2)$

(5)  $(2a + 3)(a + 4)$

(6)  $(6x - 5)(x - 1)$