

PF1.4: MOMENTUM

The momentum p of an object is the product of its mass m and velocity v , or

$$p = m \times v$$

The unit of momentum is kgms^{-1}

Momentum is a vector quantity, so it is important to remember that direction must be taken into account when doing problems on momentum.

Change in Momentum

Consider an object of mass ' m ' changing its velocity from v_i to v_f in time Δt under the action of a resultant force ΣF .

From Newton's second law of motion $\Sigma F = ma$,

$$\Sigma F = \frac{mv_f - mv_i}{\Delta t} \quad \text{since} \quad a = \frac{v_f - v_i}{\Delta t}, \quad \text{or}$$

$$\Sigma F \times \Delta t = mv_f - mv_i$$

In other words, when a resultant force ΣF acts on an object for a time duration of Δt there is a change in momentum given by $(mv_f - mv_i)$, where mv_i is the initial momentum and mv_f is the final momentum. This change in momentum is written as Δp , or

$$\Delta p = p_f - p_i = mv_f - mv_i$$

where p_i is the initial momentum and p_f is the final momentum.

Note: one must be careful when calculating Δp since change in momentum involves subtracting one vector from another.

Impulse

As shown above, when a resultant force ΣF acts on an object for a time Δt the object experiences a change in momentum. The product of this resultant force and time is called the impulse I of the force.

$$I = \Sigma F \times \Delta t = \Delta p = mv_f - mv_i$$

The unit of impulse is Ns.


It can be shown that since change in momentum is equal to impulse, then kgms^{-1} is equivalent to Ns.

Note:

1. As momentum and impulse are vectors, a sign convention in problems on momentum and impulse is essential.
2. A negative sign for the change in momentum indicates a loss of momentum; a positive sign indicates a gain in momentum.
3. Make sure you can do vector subtraction calculations.
4. Remember that ΣF is the resultant force.

Impulse is given by the area under an F- Δt graph

The area under a force-time graph gives the impulse of a force. This is very useful when the force is non-uniform (see below).



$I = F\Delta t$ is central to much of modern car design. For instance, if a test car travelling at 60km/h crashes into a wall or a large balloon (!) the change in momentum Δp for both will be the same (why?).

Since $F\Delta t = m\Delta v$ then $F\Delta t = k$ (constant)


$$\text{Hence } F \propto \frac{1}{\Delta t}$$

For the wall crash: Δt is small, therefore F is large, ie, a larger force is acting for a smaller time..

For the balloon crash: Δt is large therefore F is small, ie, a smaller force is acting for a longer time.

Conservation of Momentum

When two balls A and B collide, the action of A on B is equal and opposite to that of B on A. (Newtons 3rd Law)



Hence the rate of change of momentum of A is equal and opposite to the rate of change of momentum of B. Since the time of contact is the same for both, then the change in momentum of A is equal and opposite to the change in momentum of B.

That is, the total momentum before impact equals the total momentum after impact.

$$\begin{array}{lcl} \text{Total momentum before collision} & = & \text{total momentum after collision} \\ \Sigma p_i & = & \Sigma p_f \end{array}$$

This is known as the law of conservation of momentum. The total momentum is the same before, during and after impact, or

$$m_A u_A + m_B u_B = m_A v_A + m_B v_B$$

where m_A = mass of A, m_B = mass of B, u_A = velocity of A before impact, u_B = velocity of B before impact, v_A = velocity of A after impact, v_B = velocity of B after impact.

Note:

- Always draw a diagram when doing momentum problems.
- A sign convention (+, -) is essential (momentum and impulse are vectors). Be careful with negative and positive signs!
- If, for instance, the two cars collide and stay together after the collision, then the momentum of the two cars before the collision is equal to the momentum of the locked-together cars after the collision.
- Mathematically, problems on collisions or explosions are similar, except that for an explosion, the momentum of the system before the blast is often zero.
- In closed systems no external forces act. For example, a spacecraft colliding in deep space experiences no external force because there is no gravitational field. Systems close to the Earth's surface are systems that are not closed because the external force acting is the Earth's gravitational force. To "make" them closed one has to include the Earth as part of the system.


Momentum transfer involving the Earth

The following two situations describe how momentum is transferred to the Earth.

- A ball is thrown up: it rises against gravity and then slows down and stops, losing momentum due to the earth. The ball starts to fall under the influence of gravity, speeding up thereby giving the earth an equal and opposite momentum change. The falling ball hits the ground. The momentum is not lost. It is transferred to the earth.
- A bicycle on the flat is slowed down due to friction. The loss of momentum of the bike is transferred to the earth which gains an equal and opposite momentum.

Exercise

- 1 A tennis ball of mass 100g hits the wall horizontally at 8.0ms^{-1} east and rebounds at 6.0ms^{-1} . The contact time with the wall is 0.07s .
Calculate (a) the impulse on the ball by the wall, (b) the change in momentum of the ball
State the magnitude and direction for (a) and (b)
- 2 Referring to Question 1 above, calculate (a) the force exerted on the ball by the wall, and (b) the force exerted on the wall by the ball. State the magnitude and direction for (a) and (b)
- 3 The graph shown below shows how force varies with time for a miniature crash test dummy of mass 2 kg moving to the right. It is involved in a collision with a large concrete block set into the ground which brings it to rest. Calculate the dummy's initial speed.



- 4 A truck of mass 2500kg travelling at 20ms^{-1} west collides head on with a car of mass 800kg travelling in the opposite direction at 15ms^{-1} . The two vehicles become locked together.
(a) What is the total momentum of the two vehicles before the collision? Assume the truck's motion (west) is positive.
(b) What is the speed and direction of the car and truck immediately after the collision?
- 5 A racing car negotiating a tight bend at 30 kmh^{-1} collides with a crash barrier. The air bag in his car inflates and the time taken for it to inflate is 0.16s . The driver's head has a mass of 7.0kg .
Explain why the driver is less likely to suffer head injury in a collision with the air bag than if his head collided with the car dashboard, or other hard surface.

Answers

- 1 (a) 1.4 Ns West (b) 1.4 kgm/s West
- 2 (a) 20N West (b) 20N East
- 3 10m/s 4. (a) $3.8 \times 10^4\text{ kgm/s}$ (b) 12 m/s West
- 5
 - $F\Delta t = k$ (constant), since $m\Delta v$ is constant

$$\text{Hence, } F \propto \frac{1}{\Delta t}$$

- In other words, the force of impact is inversely proportional to the time of impact
- For the dashboard impact: Δt is small, therefore F is large, resulting in serious injury
- For the air bag impact: Δt is large, therefore F is small, resulting in a much less serious injury